

Construction of the Zero-Energy State of $SU(2)$ -Matrix Theory: Near the Origin

Jens Hoppe^{a *}, Douglas Lundholm^{a †}
and Maciej Trzetrzelewski^{a,b ‡}

^a Department of Mathematics,
Royal Institute of Technology,
KTH, 100 44 Stockholm,
Sweden

^b Institute of Physics,
Jagiellonian University,
Reymonta 4, 30-059 Kraków,
Poland

Abstract

We explicitly construct a (unique) $Spin(9) \times SU(2)$ singlet state, ϕ , involving only the fermionic degrees of freedom of the supersymmetric matrix-model corresponding to reduced 10-dimensional super Yang-Mills theory, resp. supermembranes in 11-dimensional Minkowski space. Any non-singular wavefunction annihilated by the 16 supercharges of $SU(2)$ matrix theory must, at the origin (where it is assumed to be non-vanishing) reduce to ϕ .

*e-mail: hoppe@kth.se

†e-mail: dogge@math.kth.se

‡e-mail: 33lewski@th.if.uj.edu.pl

1 Introduction

The fermionic degrees of freedom of $SU(2)$ -matrix theory (see e.g. [1]) are three $Spin(9)$ spinors, $\theta_{\hat{\alpha}A}$, ($\hat{\alpha} = 1, \dots, 16$, $A = 1, 2, 3$, $\theta_{\hat{\alpha}A}^\dagger = \theta_{\hat{\alpha}A}$), satisfying canonical anti-commutation relations

$$\{\theta_{\hat{\alpha}A}, \theta_{\hat{\beta}B}\} = \delta_{\hat{\alpha}\hat{\beta}}\delta_{AB}.$$

The corresponding $2^{8 \cdot 3}$ - dimensional Hilbert-space $\mathcal{H} = \mathcal{H}_{256} \otimes \mathcal{H}_{256} \otimes \mathcal{H}_{256}$ splits into irreducible $Spin(9)$ representations built out of the ones occurring in

$$\mathcal{H}_{256} = \mathbf{44} \oplus \mathbf{84} \oplus \mathbf{128}.$$

First determining all $Spin(9)$ singlets occurring in $\mathcal{H}$, in terms of the three representations **44**, **84**, **128** (whose elements are denoted $|st\rangle$, $|stu\rangle$ and $|t\hat{\alpha}\rangle$ respectively), the central part of the paper then is the explicit construction, out of these $Spin(9)$ singlets, of a (unique) $Spin(9) \times SU(2)$ singlet ϕ (whose relevance has been advocated by Wosiek, who was led to the existence of such a $Spin(9) \times SU(2)$ singlet using symbolic programme [2]).

In the next section we take an independent route to obtain ϕ , here proving its uniqueness, by listing all possible $Spin(7) \times SU(2)$ invariant states and then taking their (unique) linear combination such that the result is $Spin(9) \times SU(2)$ invariant.

While the three representations in $\mathcal{H}_{256}$, forming an 'Euler-triple' (cp. e.g. [3]) - quite likely relevant concerning the existence of a (unique) zero energy state for general $SU(N \geq 2)$ (note the intertwining nature of the two terms $\gamma_{\hat{\beta}\hat{\alpha}}^t \theta_{\hat{\alpha}A}$ and $\gamma_{\hat{\beta}\hat{\alpha}}^{st} \theta_{\hat{\alpha}A}$ in the supercharges of the model) - have quite a long history in supergravity theory (starting with [4]), we could not ¹ find a good reference for their Fock space representations and therefore derived them explicitly (see Appendix A) to be sure of the exact intertwining relations.

2 The construction of ϕ

According to the decomposition of $\mathcal{H}_{256} \otimes \mathcal{H}_{256}$ into irreducible representations of $Spin(9)$ (cp. e.g. [7], eq. (13)-(18), or [8]) - yielding three **44**'s and

¹at least not until September 28; our apologies to the authors of [5], [6] (whose results seem to be related to our Fock space representations given in Appendix A)

three **84**'s, which are easily seen to be (proportional to)

$$\begin{aligned} \underset{44}{|| st \rangle} &:= |su\rangle|tu\rangle + |tu\rangle|su\rangle - \frac{2}{9}\delta_{st}|uv\rangle|uv\rangle, \\ \underset{84}{|| st \rangle} &:= |suv\rangle|tuv\rangle + |tuv\rangle|suv\rangle - \frac{2}{9}\delta_{st}|uvw\rangle|uvw\rangle, \\ \underset{128}{|| st \rangle} &:= |s\hat{\alpha}\rangle|t\hat{\alpha}\rangle + |t\hat{\alpha}\rangle|s\hat{\alpha}\rangle - \frac{2}{9}\delta_{st}|u\hat{\alpha}\rangle|u\hat{\alpha}\rangle, \end{aligned}$$

and (for notational convenience we will now write α instead of $\hat{\alpha}$, in this section)

$$\begin{aligned} \underset{84}{|| stu \rangle} &:= \epsilon^{stupqrabc}|pqr\rangle|abc\rangle, \\ \underset{128}{|| stu \rangle} &:= \gamma_{\alpha\beta}^s|t\alpha\rangle|u\beta\rangle + \gamma_{\alpha\beta}^t|u\alpha\rangle|s\beta\rangle + \gamma_{\alpha\beta}^u|s\alpha\rangle|t\beta\rangle, \\ \underset{128}{|| stu \rangle}' &:= \gamma_{\alpha\beta}^{stu}|v\alpha\rangle|v\beta\rangle, \end{aligned}$$

there are 14 $Spin(9)$ singlets in $\mathcal{H}_{256} \otimes \mathcal{H}_{256} \otimes \mathcal{H}_{256}$. Nine of these involve the **128**-dimensional spinor-representations while the simplest ones are

$$\underset{44}{||| 1 \rangle} := |su\rangle_1|tu\rangle_2|st\rangle_3, \quad \underset{84}{||| 1 \rangle} := \epsilon^{stupqrabc}|stu\rangle_1|pqr\rangle_2|abc\rangle_3,$$

and the (cyclically invariant) sum of the remaining three,

$$\underset{844}{||| 1 \rangle} := |suv\rangle_1|tuv\rangle_2|st\rangle_3 + |tuv\rangle_1|st\rangle_2|suv\rangle_3 + |st\rangle_1|suv\rangle_2|tuv\rangle_3.$$

Using the 'Rarita-Schwinger' constraints (RSC) $\gamma_{\alpha\beta}^t|t\beta\rangle_A = 0$ and the intertwining relations (cp. Appendix A)

$$2\theta_{\alpha A}|st\rangle_A = \gamma_{\alpha\beta}^s|t\beta\rangle_A + \gamma_{\alpha\beta}^t|s\beta\rangle_A, \quad (1)$$

$$\theta_{\alpha A}|stu\rangle_A = \frac{i}{\sqrt{2}} \left(\gamma_{\alpha\beta}^{st}|u\beta\rangle_A + \gamma_{\alpha\beta}^{us}|t\beta\rangle_A + \gamma_{\alpha\beta}^{tu}|s\beta\rangle_A \right), \quad (2)$$

it is straightforward to calculate the action of the $SU(2)$ generators $J_A := \frac{1}{2}\epsilon_{ABC}\theta_{\alpha B}\theta_{\alpha C}$ on the above $Spin(9)$ singlets; e.g.²

$$\theta_{\alpha 1}\theta_{\alpha 2} \underset{44}{||| 1 \rangle} = \frac{13}{4}|s\beta\rangle_1|t\beta\rangle_2|st\rangle_3$$

²It may be amusing to speculate about the occurrence of the relatively large prime 13, which played a prominent role in (bosonic) string theory.

and (s.b.)

$$\theta_{\alpha 1} \theta_{\alpha 2} \underset{844}{||| 1\rangle} = -9 |s\epsilon\rangle_1 |t\epsilon\rangle_2 |st\rangle_3. \quad (3)$$

It follows that

$$\phi := \underset{44}{||| 1\rangle} + \frac{13}{36} \underset{844}{||| 1\rangle} \quad (4)$$

is $Spin(9) \times SU(2)$ invariant.

Eq. (3) easily follows when splitting the calculation into two parts:

$$-2\theta_{\alpha 1} \theta_{\alpha 2} |suv\rangle_1 |tuv\rangle_2 |st\rangle_3 =$$

$$(\gamma_{\alpha\beta}^{su} |v\beta\rangle_1 + \gamma_{\alpha\beta}^{uv} |s\beta\rangle_1 + \gamma_{\alpha\beta}^{vs} |u\beta\rangle_1)(\gamma_{\alpha\epsilon}^{tu} |v\epsilon\rangle_2 + \gamma_{\alpha\epsilon}^{uv} |t\epsilon\rangle_2 + \gamma_{\alpha\epsilon}^{vt} |u\epsilon\rangle_2) |st\rangle_3;$$

using the RSC and the (anti-)commutation relations between γ 's (in particular $\gamma^{pq} = \gamma^p \gamma^q - \delta^{pq} \mathbf{1}$ and $[\gamma^{pq}, \gamma^r] = 2\gamma^p \delta^{qr} - 2\gamma^q \delta^{pr}$), each of the nine terms becomes proportional to $|s\epsilon\rangle_1 |t\epsilon\rangle_2 |st\rangle_3$, the respective coefficients being: 0 for $\gamma^{su} \gamma^{tu}$, $\gamma^{vs} \gamma^{vt}$; 72 for $\gamma^{uv} \gamma^{uv}$; 3 for $\gamma^{su} \gamma^{vt}$, $\gamma^{vs} \gamma^{tu}$; -15 for each of the remaining four $\gamma^{su} \gamma^{uv}$, $\gamma^{uv} \gamma^{tu}$, $\gamma^{uv} \gamma^{vt}$, $\gamma^{vs} \gamma^{uv}$.

Concerning the second part in (3), one notes that

$$-\sqrt{8}i\theta_{\alpha 1} \theta_{\alpha 2} (|tuv\rangle_1 |st\rangle_2 + |st\rangle_1 |tuv\rangle_2) |suv\rangle_3 =$$

$$(\gamma_{\alpha\beta}^{tu} |v\beta\rangle_1 + \gamma_{\alpha\beta}^{uv} |t\beta\rangle_1 + \gamma_{\alpha\beta}^{vt} |u\beta\rangle_1)(\gamma_{\alpha\epsilon}^s |t\epsilon\rangle_2 + \gamma_{\alpha\epsilon}^t |s\epsilon\rangle_2) |suv\rangle_3 \\ + (\gamma_{\alpha\beta}^s |t\beta\rangle_1 + \gamma_{\alpha\beta}^t |s\beta\rangle_1)(\gamma_{\alpha\epsilon}^{tu} |v\epsilon\rangle_2 + \gamma_{\alpha\epsilon}^{uv} |t\epsilon\rangle_2 + \gamma_{\alpha\epsilon}^{vt} |u\epsilon\rangle_2) |suv\rangle_3,$$

which gives rise to 12 terms, 6 of which cancel in pairs ($\gamma^{uv} \gamma^s$, $\gamma^{tu} \gamma^t$ and $\gamma^{vt} \gamma^t$), due to antisymmetry of γ^{su} (as a matrix) and $|suv\rangle$, while (using again $\gamma^{pq} = \gamma^p \gamma^q - \delta^{pq} \mathbf{1}$ and the transformation of the γ^w 's as a vector when commuting with γ^{pq}) the 3 remaining terms containing $|t\epsilon\rangle_2$, resulting in

$$5(\gamma_{\beta\epsilon}^v |s\beta\rangle_1 |u\epsilon\rangle_2 - \gamma_{\beta\epsilon}^u |s\beta\rangle_1 |v\epsilon\rangle_2) |suv\rangle_3,$$

are cancelled by those arising from the terms containing $|t\beta\rangle_1$.

Perhaps it is worth noting that the image (under the action of any of the $SU(2)$ generators, say J_3) of $\underset{84}{||| 1\rangle}$ (which is *not* needed for the $Spin(9) \times$

$SU(2)$ singlet) is, using $\gamma^{stpq} \propto \gamma^{xyzvw} \epsilon_{xyzvwstpq}$, proportional to $\gamma_{\beta\epsilon}^s |u\beta\rangle_1 |v\epsilon\rangle_2 |suv\rangle_3$ (and that the apparent 'puzzle' of the $Spin(9)$ singlet $\gamma_{\beta\epsilon}^{su} |t\beta\rangle_1 |t\epsilon\rangle_2 |suv\rangle_3$ *not* entering these considerations is 'resolved' by observing that only the cyclically invariant combination $\underset{844}{||| 1\rangle}$ was considered - in which $\gamma_{\beta\epsilon}^{su}$ terms appear *symmetrized* in β and ϵ , i.e. cancelling each other).

3 The construction of ϕ out of $Spin(7) \times SU(2)$ singlets

Here we reproduce the result obtained in the previous section using an independent approach. Our strategy is to take advantage of the natural $Spin(7)$ covariance of fermionic creation operators $\lambda_{\alpha A}$, $\alpha = 1, \dots, 8$, corresponding to the full $Spin(9) \times SU(2)$ model. To do so we first write the fermionic $Spin(9)$ generators $M_{st} = \frac{1}{4} \gamma_{\hat{\alpha}\hat{\beta}}^{st} \theta_{\hat{\alpha}A} \theta_{\hat{\beta}A}$ as $(i, j = 1, \dots, 7)$

$$M_{ij} = \frac{1}{2} \Gamma_{\alpha\beta}^{ij} \lambda_{\alpha A} \lambda_{\beta A}^\dagger, \quad M_{j8} = \frac{i}{4} \Gamma_{\alpha\beta}^j (\lambda_{\alpha A} \lambda_{\beta A} + \lambda_{\alpha A}^\dagger \lambda_{\beta A}^\dagger),$$

$$M_{89} = -\frac{i}{2} (\lambda_{\alpha A} \lambda_{\alpha A}^\dagger - 12) \quad M_{j9} = -\frac{1}{4} \Gamma_{\alpha\beta}^j (\lambda_{\alpha A} \lambda_{\beta A} - \lambda_{\alpha A}^\dagger \lambda_{\beta A}^\dagger),$$

where we use the conventions for γ^s as given in Appendix A.

The condition $M_{ij}\phi = 0$ is the $Spin(7)$ invariance of ϕ while $M_{89}\phi = 0$ tells us that $\phi \in \mathcal{F}_{12}$ (where $\mathcal{F}_{n_F}$ denotes the sector with n_F fermions). Therefore we are led to search for a combination of $Spin(7) \times SU(2)$ invariant states in $\mathcal{F}_{12}$ such that

$$\Gamma_{\alpha\beta}^i \lambda_{\alpha A} \lambda_{\beta A} \phi = \Gamma_{\alpha\beta}^i \lambda_{\alpha A}^\dagger \lambda_{\beta A}^\dagger \phi = 0. \quad (5)$$

3.1 The number of $Spin(7) \times SU(2)$ invariant states

In order to solve Eqn. (5) we attempt to list all $Spin(7) \times SU(2)$ invariant states in $\mathcal{F}_{12}$. In doing so it is helpful to first calculate the number D_{n_F} of such states appearing in $\mathcal{F}_{n_F}$ for arbitrary n_F . This is done following the lines of [9] by writing $\mathcal{F}_{n_F}$ as

$$\mathcal{F}_{n_F} = Alt(\otimes_{l=1}^{n_F} F_l^{s=1/2}) = \mathcal{F}_{n_F, j=0} \oplus \mathcal{F}_{n_F, j=1/2} \oplus \dots,$$

where $F_l^{s=1/2}$ is a vector space spanned by $\lambda_l^{s=1/2}|0\rangle$ (operators $\lambda_l^{s=1/2}$ are assumed to carry spin $s = 1/2$ of $SO(7)$) and where $\mathcal{F}_{n_F, j}$ is $\mathcal{F}_{n_F}$ projected into subspaces with given $SO(7)$ angular momentum. Therefore the dimensions of subspaces with angular momentum $j = 0$ are

$$D_{n_F}^{Spin(7) \times SU(2)} = \int d\mu_{SO(7)} \underbrace{\chi^{SO(7)}_{j=0}}_{=1} \int d\mu_{SU(2)} \chi_{Alt}^{[n_F]}(R),$$

where $d\mu_{SO(7)}$ and $d\mu_{SU(2)}$ are $SO(7)$ and $SU(2)$ invariant measures, R is the adjoint representation of $SU(2)$ and $j = 1/2$ representation of $SO(7)$, i.e. $R = R^{SO(7), j=1/2} \otimes R^{SU(2), j=1}$. The characters χ can be read off directly from the Weyl character formula while the antisymmetric power of $\chi(R)$ is given by the Frobenius formula (see e.g [10])

$$\chi_{Alt}^{[n_F]}(R) = \sum_{\sum_k k i_k = n_F} (-1)^{\sum_k i_k} \prod_{k=1}^{n_B} \frac{1}{i_k!} \frac{\chi^{i_k}(R^k)}{k^{i_k}},$$

(here R is considered as a matrix). Taking all into consideration we find that the generating function for the numbers D_{n_F} is

$$\sum_{n_F=0}^{24} D_{n_F} b^{n_F} = 1 + 2b^4 + 5b^8 + 7b^{12} + 5b^{16} + 2b^{20} + b^{24}.$$

Note the duality between $\mathcal{F}_{n_F}$ and $\mathcal{F}_{24-n_F}$, i.e. the particle-hole symmetry.

3.2 The construction of $Spin(7) \times SU(2)$ invariant states and ϕ

We now proceed to construct the $Spin(7) \times SU(2)$ invariant states in $\mathcal{F}_4$, $\mathcal{F}_8$ and finally in $\mathcal{F}_{12}$ (note that there are no such states in $\mathcal{F}_{4n+2}$, $n = 0, \dots, 5$ and $\mathcal{F}_{2n+1}$, $n = 0, \dots, 11$). Let us first consider operators

$$b_{AB} := \lambda_{\alpha A} \lambda_{\alpha B}, \quad b_{AB}^i := \Gamma_{\alpha\beta}^i \lambda_{\alpha A} \lambda_{\beta B}.$$

In the $\mathcal{F}_4$ sector there are only two such states for which we choose

$$v_1|0\rangle, \quad v_2|0\rangle \quad v_1 := b_{AB} b_{AB} \quad v_2 := b_{AB}^i b_{AB}^i.$$

They are not orthogonal as the overlap matrix $G = [\langle 0|v_i^\dagger v_j|0\rangle]_{i,j=1,2}$ is (see Appendix B)

$$G = \begin{pmatrix} 1344 & 2688 \\ 2688 & 45696 \end{pmatrix}, \quad \det G \neq 0. \quad (6)$$

We developed a symbolic programme written in Mathematica to confirm that indeed G is of this form.

In the $\mathcal{F}_8$ fermion sector there are 5 invariant states. Three of them are simply

$$v_1 v_1|0\rangle, \quad v_1 v_2|0\rangle, \quad v_2 v_2|0\rangle.$$

The remaining two are e.g.

$$w_1|0\rangle, \quad w_2|0\rangle, \quad w_1 = b_{AB}^i b_{BC}^i b_{CD}^j b_{DA}^j, \quad w_2 = b_{AB}^i b_{BC}^j b_{CD}^i b_{DA}^j.$$

We checked in Mathematica that these states are linearly independent.

Finally in the $\mathcal{F}_{12}$ sector we should have 7 states. Considering the previous sectors we can construct already 8. They are

$$\begin{aligned} &v_1 v_1 v_1|0\rangle, \quad v_1 v_1 v_2|0\rangle, \quad v_1 v_2 v_2|0\rangle, \quad v_2 v_2 v_2|0\rangle, \\ &v_1 w_1|0\rangle, \quad v_2 w_1|0\rangle, \quad v_1 w_2|0\rangle, \quad v_2 w_2|0\rangle. \end{aligned}$$

Accordingly there should be one relation between them. Indeed, we found that

$$\begin{aligned} &4358v_1 v_1 v_1|0\rangle + 2652v_1 v_1 v_2|0\rangle + 984v_1 v_2 v_2|0\rangle + 63v_2 v_2 v_2|0\rangle - 528v_1 w_1|0\rangle \\ &- 88v_1 w_2|0\rangle + 24v_2 w_1|0\rangle - 152v_2 w_2|0\rangle = 0, \end{aligned}$$

and that there are no other identities among these 8 states. Therefore we can choose the $\mathcal{F}_{12}$ basis to be e.g.

$$\begin{aligned} &r_1 = v_1 v_1 v_1|0\rangle, \quad r_2 = v_1 v_1 v_2|0\rangle, \quad r_3 = v_1 v_2 v_2|0\rangle, \quad r_4 = v_2 v_2 v_2|0\rangle, \\ &r_5 = v_1 w_1|0\rangle, \quad r_6 = v_2 w_1|0\rangle, \quad r_7 = v_1 w_2|0\rangle. \end{aligned}$$

Finally we checked that there exists a unique combination of r_i such that Eqn. (5) is satisfied. The result reads

$$\chi = 326304r_1 + 488136r_2 + 72612r_3 + 1377r_4 + 114576r_5 - 176528r_6 + 10296r_7,$$

and is proportional to (4).

4 Outlook

Having solved a (physically relevant) representation theoretic question, let us now make a comment on the problem of determining the full zero-energy eigenfunction Ψ of the Hamiltonian

$$H = -\Delta + \frac{1}{2}(\epsilon_{ABC}x_s B x_{tC})^2 + i x_{sC} \epsilon_{ABC} \gamma_{\hat{\alpha}\hat{\beta}}^s \theta_{\hat{\alpha}A} \theta_{\hat{\beta}B},$$

which on the physical space of $SU(2)$ invariant states is equal to the square of each of the supercharges

$$\mathcal{Q}_{\hat{\beta}} = -i\partial_{sA}\gamma_{\hat{\beta}\hat{\alpha}}^s\theta_{\hat{\alpha}A} + \frac{1}{2}\epsilon_{ABC}x_{sB}x_{tC}\gamma_{\hat{\beta}\hat{\alpha}}^{st}\theta_{\hat{\alpha}A} = D_{\hat{\beta}} + V_{\hat{\beta}}.$$

Due to elliptic regularity (see e.g. [11]), any solution to $H\Psi = 0$ must be smooth. Accordingly, one can, around the origin, write Ψ in terms of a power series in the coordinates,

$$\Psi(x) = \sum_{k=0}^N \Psi^{(k)}(x) = \psi^{(0)} + x_{tA}\psi_{tA}^{(1)} + \frac{1}{2}x_{tA}x_{uB}\psi_{tA,uB}^{(2)} + \dots + \Psi^{(N)}(x),$$

with $\psi_{t_1A_1\dots t_kA_k}^{(k)} \in \mathcal{H}$, and $\Psi^{(k)}$ vanishing to order k at $x = 0$.

Examining the equations $\mathcal{Q}_{\hat{\beta}}\Psi = 0$ to each order in the coordinates, we find

$$D_{\hat{\beta}}\Psi^{(1)} = 0, \quad D_{\hat{\beta}}\Psi^{(2)} = 0, \quad D_{\hat{\beta}}\Psi^{(k+3)} + V_{\hat{\beta}}\Psi^{(k)} = 0, \quad k = 0, 1, 2, \dots$$

i.e.

$$\gamma_{\hat{\beta}\hat{\alpha}}^t\theta_{\hat{\alpha}A}\psi_{tA}^{(1)} = 0,$$

$$\gamma_{\hat{\beta}\hat{\alpha}}^t\theta_{\hat{\alpha}A}\psi_{tA,uB}^{(2)} = 0,$$

$$\gamma_{\hat{\beta}\hat{\alpha}}^t\theta_{\hat{\alpha}A}\psi_{tA,uB,vC}^{(3)} + \frac{1}{2}\epsilon_{ABC}\gamma_{\hat{\beta}\hat{\alpha}}^{uv}\theta_{\hat{\alpha}A}\psi^{(0)} = 0,$$

etc. for all $\hat{\beta}$. Note the three separate towers of equations relating $\Psi^{(k)}$ to $\Psi^{(k+3)}$ via intertwiners.

In any case, using that Ψ must be $Spin(9)$ invariant [12], one concludes that $\psi^{(0)}$ must be a scalar multiple of the state we constructed in this paper.

Acknowledgments We would like to thank V. Bach, G. M. Graf, R. Suter and J. Wosiek for discussions, as well as the Swedish Research Council and the Marie Curie Training Network ENIGMA (contract MRNT-CT-2004-5652) for financial support.

Appendix A

In this Appendix we work with a single $\mathcal{H}_{256}$. A Fock space representation of $\mathcal{H}_{256}$ can be obtained by introducing fermionic creation operators λ_α and annihilation operators $\frac{\partial}{\partial \lambda_\alpha} = \lambda_\alpha^\dagger$ via

$$\theta_\alpha = \frac{1}{\sqrt{2}}(\lambda_\alpha + \lambda_\alpha^\dagger), \quad \theta_{\alpha+8} = \frac{1}{i\sqrt{2}}(\lambda_\alpha - \lambda_\alpha^\dagger).$$

A basis of the Hilbert space $\mathcal{H}_{256}$ is obtained by acting with products of the λ_α 's on the fermion vacuum state $|0\rangle$ defined by $\lambda_\alpha^\dagger |0\rangle = 0$. The $Spin(9)$ generators $M_{st} = \frac{1}{4}\gamma_{\hat{\alpha}\hat{\beta}}^{st}\theta_{\hat{\alpha}}\theta_{\hat{\beta}}$, $s, t = 1, \dots, 9$ where $\gamma^{st} = \frac{1}{2}[\gamma^s, \gamma^t]$ and γ^s are 16×16 , real, symmetric matrices satisfying $\{\gamma^s, \gamma^t\} = 2\delta^{st}\mathbf{1}_{16 \times 16}$, then become

$$\begin{aligned} M_{ij} &= \frac{1}{2}\Gamma_{\alpha\beta}^{ij}\lambda_\alpha\lambda_\beta^\dagger, & M_{j8} &= \frac{i}{4}\Gamma_{\alpha\beta}^j(\lambda_\alpha\lambda_\beta + \lambda_\alpha^\dagger\lambda_\beta^\dagger), \\ M_{89} &= -\frac{i}{2}(\lambda_\alpha\lambda_\alpha^\dagger - 4) & M_{j9} &= -\frac{1}{4}\Gamma_{\alpha\beta}^j(\lambda_\alpha\lambda_\beta - \lambda_\alpha^\dagger\lambda_\beta^\dagger), \end{aligned} \quad (7)$$

when choosing

$$\gamma^j = \begin{bmatrix} 0 & i\Gamma^j \\ -i\Gamma^j & 0 \end{bmatrix}, \quad \gamma^8 = \begin{bmatrix} 0 & \mathbf{1}_{8 \times 8} \\ \mathbf{1}_{8 \times 8} & 0 \end{bmatrix}, \quad \gamma^9 = \begin{bmatrix} \mathbf{1}_{8 \times 8} & 0 \\ 0 & -\mathbf{1}_{8 \times 8} \end{bmatrix},$$

with Γ^i being 8×8 , purely imaginary, antisymmetric matrices satisfying $\{\Gamma^i, \Gamma^j\} = 2\delta^{ij}\mathbf{1}_{8 \times 8}$ ³

As already mentioned, the Hilbert space $\mathcal{H}_{256}$ decomposes into three irreducible representations whose elements will be denoted by $|st\rangle$, $|stu\rangle$ and $|s\hat{\alpha}\rangle$ respectively.

44

An explicit presentation of the **44** in terms of creation operators λ_α was given in [13] as follows:

$$|i \neq j\rangle = b_i b_j |0\rangle, \quad |jj\rangle = \left(b_j^2 - \frac{1}{9}\mathbf{b}^2\right) |0\rangle,$$

³Furthermore one may choose $i\Gamma_{\alpha 8}^j = \delta_\alpha^j$, $i\Gamma_{kl}^j = -c_{jkl}$ with totally antisymmetric octonionic structure constants $c_{ijk} = +1$ for $(ijk) = (123), (165), (246), (435), (147), (367), (257)$.

$$\begin{aligned}
|j8\rangle &= \frac{1}{2}b_j \left(1 - \frac{2}{9}\mathbf{b}^2\right) |0\rangle, & |j9\rangle &= -\frac{i}{2}b_j \left(1 + \frac{2}{9}\mathbf{b}^2\right) |0\rangle, \\
|88\rangle &= \frac{1}{2} \left(|0\rangle - \frac{2}{9}\mathbf{b}^2|0\rangle + |8\rangle\right), & |99\rangle &= -\frac{1}{2} \left(|0\rangle + \frac{2}{9}\mathbf{b}^2|0\rangle + |8\rangle\right), \\
|89\rangle &= -\frac{i}{2} (|0\rangle - |8\rangle), & &
\end{aligned} \tag{8}$$

where

$$b_j := \frac{i}{4}\Gamma_{\alpha\beta}^j \lambda_\alpha \lambda_\beta, \quad \mathbf{b}^2 := \sum_{i=1}^7 b_i b_i, \quad |8\rangle := \lambda_1 \dots \lambda_8 |0\rangle.$$

While it is convenient to work with the overcomplete set of states $|st\rangle = |ts\rangle$, satisfying $\sum_{s=1}^9 |ss\rangle = 0$ and transforming according to

$$M_{st}|uv\rangle = \delta_{tu}|sv\rangle - \delta_{su}|tv\rangle + \delta_{tv}|su\rangle - \delta_{sv}|tu\rangle, \tag{9}$$

one should be aware of the fact that they are *not* orthonormal; rather

$$\langle st|s't'\rangle = \frac{1}{2}(\delta_{ss'}\delta_{tt'} + \delta_{st'}\delta_{s't})(1 - \delta_{st})(1 - \delta_{s't'}) + \delta_{st}\delta_{s't'} \left(\delta_{ss'} - \frac{1}{9}\right)$$

(in accordance with $|s \neq t\rangle \cong \frac{1}{2}(|st\rangle + |ts\rangle)$, $|tt\rangle \cong \frac{1}{2}(|tt\rangle + |tt\rangle) - \frac{1}{9} \sum_u |uu\rangle$, where $|st\rangle$ are unconstrained tensor product-states satisfying $\langle st|s't'\rangle = \delta_{ss'}\delta_{tt'}$)

(8) follows from (9) when starting with the 27-dimensional traceless symmetric $U(1)$ ($\cong M_{89}$)-invariant $Spin(7)$ representation containing $|i \neq j\rangle = b_i b_j |0\rangle$:

$$|9j\rangle = M_{9k}|jk\rangle = -i(b_k + b_k^\dagger)b_j b_k |0\rangle = -\frac{i}{2}(b_j + 2b_k^2 b_j)|0\rangle,$$

$$|8j\rangle = M_{8k}|jk\rangle = \frac{1}{2}(b_j - 2b_k^2 b_j)|0\rangle,$$

$$M_{9j}|j9\rangle = |99\rangle - |jj\rangle = -\left(\frac{1}{2} + b_j^2 + b_j^2 b_k^2\right)|0\rangle,$$

$$M_{8j}|j8\rangle = |88\rangle - |jj\rangle = \left(\frac{1}{2} - b_j^2 + b_j^2 b_k^2\right)|0\rangle,$$

(with $j \neq k$, no sum) implying

$$|88\rangle - |99\rangle = (1 + 2b_j^2 b_k^2)|0\rangle = |0\rangle + |8\rangle,$$

$$7(|88\rangle + |99\rangle) = -2(|88\rangle + |99\rangle + \mathbf{b}^2|0\rangle),$$

i.e.

$$|88\rangle + |99\rangle = -\frac{2}{9}\mathbf{b}^2|0\rangle,$$

hence

$$|jj\rangle = b_j^2|0\rangle - \frac{1}{9}\mathbf{b}^2|0\rangle.$$

Note that one may use

$$\begin{aligned} [b_i, b_j^\dagger] &= \frac{1}{2}M_{ij} + \delta_{ij} \left(1 - \frac{1}{4}\lambda_\alpha \lambda_\alpha^\dagger\right), \\ [M_{ij}, b_k] &= \delta_{jk}b_i - \delta_{ik}b_j, \end{aligned} \tag{10}$$

which follows from

$$\begin{aligned} [Tr(\lambda A \lambda), Tr(\lambda^\dagger B \lambda^\dagger)] &= -4Tr(\lambda B A \lambda^\dagger) + 2Tr(AB), \\ [Tr(\lambda A \lambda^\dagger), Tr(\lambda^\dagger B \lambda^\dagger)] &= 2Tr(\lambda^\dagger A B \lambda^\dagger). \end{aligned} \tag{11}$$

Consistency conditions such as ($i \neq j$)

$$|j8\rangle = M_{8i}|ij\rangle = \frac{1}{2}M_{j8}|88\rangle = M_{j9}|98\rangle,$$

lead to useful (Fierz-)identities ($j \neq k$)

$$2b_j b_k^2|0\rangle = \frac{2}{3}b_j^3|0\rangle = \frac{2}{9}b_j \mathbf{b}^2|0\rangle = b_j^\dagger|8\rangle.$$

84

The construction of states $|stu\rangle$ transforming according to the antisymmetric representation **84** can be done analogously, starting with

$$|ijk\rangle := \sqrt{\frac{2}{9}}(b_i b_{jk} + b_k b_{ij} + b_j b_{ki})|0\rangle,$$

where $b_{jk} := \frac{1}{4}Tr(\lambda \Gamma^{jk} \lambda)$. In proving

$$\langle ijk|i'j'k'\rangle = \delta_{ii'}\delta_{jj'}\delta_{kk'}, \quad i < j < k, \quad i' < j' < k',$$

it is helpful to use (10) and the commutation rules

$$[M_{ij}, b_{kl}] = \delta_{jk}b_{il} + \delta_{il}b_{jk} - \delta_{ik}b_{jl} - \delta_{jl}b_{ik},$$

$$[b_k^\dagger, b_{ij}] = \frac{i}{4} \text{Tr}(\lambda \Gamma^{ij} \Gamma^k \lambda^\dagger), \quad \langle 0 | b_{ij}^\dagger b_{kl} | 0 \rangle = \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}$$

which follow from (11).

All other states of the **84** can be obtained by application of the M_{st} operators on $|ijk\rangle$ using

$$M_{st}|pqr\rangle = \delta_{tp}|sqr\rangle - \delta_{tq}|spr\rangle + \delta_{tr}|spq\rangle - \delta_{sp}|tqr\rangle + \delta_{sq}|tpr\rangle - \delta_{sr}|tpq\rangle,$$

from which it follows that for fixed i, j and k we have

$$|ij8\rangle = M_{8k}|ijk\rangle, \quad |ij9\rangle = M_{9k}|ijk\rangle,$$

$$|i89\rangle = -M_{9j}|ij8\rangle.$$

Again, using the independence of k in the above formulas resp. additional relations among various states defined via b_i and b_{jk} one can show that

$$|ij8\rangle = \frac{1}{\sqrt{2}} (b_{ij}|0\rangle + b_{ij}^\dagger|8\rangle), \quad |ij9\rangle = -\frac{i}{\sqrt{2}} (b_{ij}|0\rangle - b_{ij}^\dagger|8\rangle),$$

$$\begin{aligned} |i89\rangle &= -M_{9j}M_{8k}|ijk\rangle = i(b_j + b_j^\dagger)(b_k - b_k^\dagger)|ijk\rangle = \\ &= -i(b_j b_k^\dagger - b_j b_k + b_j^\dagger b_k^\dagger - b_j^\dagger b_k)|ijk\rangle; \end{aligned}$$

out of the four terms the second and third one each gives zero (due to being independent of $j \neq k$, while the sum over j and k vanishes), while the fourth ($= i(-b_k b_j^\dagger - \frac{1}{2}M_{jk})|ijk\rangle = -ib_k b_j^\dagger|ijk\rangle$) equals the first (with j and k interchanged!) - which is easily calculated to give $\sqrt{\frac{2}{9}} \frac{3}{2} i b_j b_{ij} |0\rangle$; so

$$|i89\rangle = \frac{i}{\sqrt{2}} (b_j b_{ij} + b_k b_{ik}) |0\rangle = \frac{i}{3\sqrt{2}} \sum_{l=1}^7 b_l b_{il} |0\rangle \quad i \neq j \neq k \neq i,$$

from which it follows that $\Gamma_{[\alpha\beta]}^{ij} \Gamma_{[\rho\epsilon]}^j + \Gamma_{[\alpha\beta]}^{ik} \Gamma_{[\rho\epsilon]}^k$ (no sum, $i \neq j \neq k \neq i$) must be independent of j and k (true even if $j = k$).

128

The **128** representation comprises all odd fermion states in $\mathcal{H}_{256}$. As a convenient definition one may take

$$|t\hat{\alpha}\rangle := \frac{2}{11} \gamma_{\hat{\alpha}\hat{\beta}}^s \theta_{\hat{\beta}} |st\rangle,$$

which does transform according to

$$M_{uv}|t\hat{\alpha}\rangle = \delta_{vt}|u\hat{\alpha}\rangle - \delta_{ut}|v\hat{\alpha}\rangle - \frac{1}{2}\gamma_{\hat{\alpha}\hat{\beta}}^{uv}|t\hat{\beta}\rangle$$

and explicitly exhibits the crucial RSC $\gamma_{\hat{\alpha}\hat{\beta}}^t|t\hat{\alpha}\rangle = 0$. The intertwining relation

$$2\theta_{\hat{\alpha}}|st\rangle = \gamma_{\hat{\alpha}\hat{\beta}}^s|t\hat{\beta}\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^t|s\hat{\beta}\rangle \quad (12)$$

follows when using

$$\gamma_{\hat{\alpha}\hat{\beta}}^{su}\theta_{\hat{\beta}}|tu\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^{tu}\theta_{\hat{\beta}}|su\rangle = 9\theta_{\hat{\alpha}}|st\rangle;$$

it is also true that

$$\theta_{\hat{\alpha}}|stu\rangle = \frac{i}{\sqrt{2}} \left(\gamma_{\hat{\alpha}\hat{\beta}}^{st}|u\hat{\beta}\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^{us}|t\hat{\beta}\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^{tu}|s\hat{\beta}\rangle \right), \quad (13)$$

respectively

$$|t\hat{\alpha}\rangle = \frac{i\sqrt{2}}{42}\gamma_{\hat{\alpha}\hat{\beta}}^{sv}\theta_{\hat{\beta}}|svt\rangle,$$

-which of course could have alternatively been used to define $|t\hat{\alpha}\rangle$.

Intertwiners

The above intertwining relations (12) and (13) as well as the ones below (explicitly checked on the computer), we believe to be crucial for the construction of the full zero energy state;

$$\gamma_{\hat{\alpha}\hat{\beta}}^{stu}\theta_{\hat{\beta}}|stu\rangle = 0,$$

$$|t\hat{\alpha}\rangle = \gamma_{\hat{\alpha}\hat{\beta}}^t\theta_{\hat{\beta}}|tt\rangle,$$

$$|stu\rangle = \frac{i}{44\sqrt{2}} \left(\theta\gamma^{st}\theta|uv\rangle + \theta\gamma^{tuv}\theta|sv\rangle + \theta\gamma^{tuv}\theta|sv\rangle \right),$$

$$|st\rangle = \frac{i}{168\sqrt{2}} \left(\theta\gamma^{su}\theta|tuv\rangle + \theta\gamma^{tuv}\theta|su\rangle \right),$$

$$\gamma_{\hat{\alpha}\hat{\beta}}^{su}\theta_{\hat{\beta}}|tu\rangle - \gamma_{\hat{\alpha}\hat{\beta}}^{tu}\theta_{\hat{\beta}}|su\rangle = \frac{11}{6\sqrt{2}i}\gamma_{\hat{\alpha}\hat{\beta}}^u\theta_{\hat{\beta}}|stu\rangle,$$

$$\gamma_{\hat{\alpha}\hat{\beta}}^{sv}\theta_{\hat{\beta}}|tuv\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^{uv}\theta_{\hat{\beta}}|stv\rangle + \gamma_{\hat{\alpha}\hat{\beta}}^{tv}\theta_{\hat{\beta}}|usv\rangle = 9\theta_{\hat{\alpha}}|stu\rangle,$$

$$\gamma_{\hat{\alpha}\hat{\beta}}^s\theta_{\hat{\beta}}|su\rangle = \frac{11i\sqrt{2}}{84}\gamma_{\hat{\alpha}\hat{\beta}}^{st}\theta_{\hat{\beta}}|stu\rangle.$$

Appendix B

In this Appendix we derive the form of the Gram matrix (6) related to the $\mathcal{F}_4$ sector. A similar procedure can be applied for the other sectors.

We will use the following notation

$$\begin{aligned} (\lambda_A A \lambda_B) &:= \lambda_{A\alpha} A^{\alpha\beta} \lambda_{B\beta}, & (\lambda_A A \lambda_B^\dagger) &:= \lambda_{A\alpha} A^{\alpha\beta} \lambda_{B\beta}^\dagger, \\ b_{AB} &:= (\lambda_A \mathbf{1} \lambda_B), & b_{AB}^i &:= (\lambda_A \Gamma^i \lambda_B), & b_{AB}^{ij} &:= (\lambda_A \Gamma^{ij} \lambda_B), \\ M_{AB} &:= (\lambda_A \mathbf{1} \lambda_B^\dagger), & M_{AB}^i &:= (\lambda_A \Gamma^i \lambda_B^\dagger), & M_{AB}^{ij} &:= (\lambda_A \Gamma^{ij} \lambda_B^\dagger). \end{aligned}$$

It is now useful to write down the generalization of commutation relations (11) for operators involving $\lambda_{\alpha A}$ and $\lambda_{\alpha A}^\dagger$ with color indices. We have

$$\begin{aligned} [(\lambda_A A \lambda_B), (\lambda_C^\dagger B \lambda_D^\dagger)] &= -(\lambda_C B A \lambda_B^\dagger) \delta_{AD} - (\lambda_D B^T A^T \lambda_A^\dagger) \delta_{BC} \\ &+ (\lambda_D B^T A \lambda_B^\dagger) \delta_{AC} + (\lambda_C B A^T \lambda_A^\dagger) \delta_{BD} - (AB^T) \delta_{AC} \delta_{BD} + (AB) \delta_{AD} \delta_{BC}, \\ [(\lambda_A A \lambda_B^\dagger), (\lambda_C B \lambda_D)] &= (\lambda_A A B \lambda_D) \delta_{BC} - (\lambda_A A B^T \lambda_C) \delta_{BD}. \end{aligned}$$

The commutators of the $b - b$ type are now

$$\begin{aligned} [b_{AB}^\dagger, b_{CD}] &= M_{CB} \delta_{DA} + M_{DA} \delta_{CB} - M_{DB} \delta_{CA} - M_{CA} \delta_{BD} + 8(\delta_{AC} \delta_{BD} - \delta_{AD} \delta_{BC}), \\ [b_{AB}^\dagger, b_{CD}^i] &= M_{CB}^i \delta_{AD} - M_{DA}^i \delta_{BC} + M_{DB}^i \delta_{AC} - M_{CA}^i \delta_{BD}, \\ [b_{AB}^{\dagger i}, b_{CD}] &= -M_{CB}^i \delta_{AD} + M_{DA}^i \delta_{BC} + M_{DB}^i \delta_{AC} - M_{CA}^i \delta_{BD}, \\ [b_{AB}^{\dagger i}, b_{CD}^j] &= M_{CB}^{ij} \delta_{AD} + M_{DA}^{ij} \delta_{CB} + M_{DB}^{ij} \delta_{CA} + M_{CA}^{ij} \delta_{DB} \\ &- \delta^{ij} (M_{CB} \delta_{AD} + M_{DA} \delta_{CB} + M_{DB} \delta_{CA} + M_{CA} \delta_{DB}) + 8\delta^{ij} (\delta_{AC} \delta_{BD} + \delta_{AD} \delta_{BC}), \end{aligned}$$

while the commutators of the $M - b$ type are

$$\begin{aligned} [M_{AB}, b_{CD}] &= b_{AD} \delta_{BC} - b_{AC} \delta_{BD}, \\ [M_{AB}, b_{CD}^i] &= b_{AD}^i \delta_{BC} + b_{AC}^i \delta_{BD}, \\ [M_{AB}^i, b_{CD}] &= b_{AD}^i \delta_{BC} - b_{AC}^i \delta_{BD}, \\ [M_{AB}^i, b_{CD}^j] &= b_{AD}^{ij} \delta_{BC} + b_{AC}^{ij} \delta_{BD} + b_{AD} \delta_{BC} + b_{AC} \delta_{BD}. \end{aligned}$$

Now it is straightforward to evaluate the scalar product $\langle 0 | v_1^\dagger v_1 | 0 \rangle$ with $v_1 := b_{AB} b_{AB}$, we have

$$[b_{AB}^\dagger, b_{CD} b_{CD}] = -4b_{AD} M_{DB} + 4b_{BD} M_{DA} + 28b_{AB},$$

which implies

$$[v_1^\dagger, v_1] = 8M_{AB}M_{AB} - 128M_{AA} + 16b_{AB}M_{BC}b_{CA}^\dagger + 40b_{AB}b_{AB}^\dagger + 1344,$$

hence $\langle 0|v_1^\dagger v_1|0\rangle = 1344$.

To calculate the scalar product $\langle 0|v_2^\dagger v_1|0\rangle$ with $v_2 := b_{AA}^i b_{BB}^i$ we need

$$[b_{AB}^{i\dagger}, b_{CD}b_{CD}] = -4b_{AC}M_{CB}^i + 4b_{BC}M_{CA}^i - 4b_{AB}^i + 4\delta_{AB}b_{CC}^i,$$

which gives

$$[b_{AA}^{i\dagger}, b_{CD}b_{CD}] = 8b_{AA}^i, \quad [v_2^\dagger, v_1] = 16b_{AB}^i b_{AB}^{i\dagger} - 224M_{AA} + 2688,$$

hence $\langle 0|v_2^\dagger v_1|0\rangle = 2688$

Finally the scalar product $\langle 0|v_2^\dagger v_2|0\rangle$ is obtained with use of

$$[b_{AA}^{i\dagger}, b_{BB}^j b_{CC}^j] = 8(b_{AA}^j M_{BB}^{ij} - b_{AA}^i M_{BB}^j) + 136b_{AA}^i,$$

hence

$$\begin{aligned} [v_2^\dagger, v_2] &= 16b_{AA}^i (M_{BB}^{ij} - \delta^{ij} M_{BB}) b_{CC}^{j\dagger} \\ &+ 32M_{AA}^{ij} M_{BB}^{ij} + 224M_{AA}M_{BB} + 1120M_{AA} + 352b_{AA}^i b_{BB}^{i\dagger} + 45696, \end{aligned}$$

therefore $\langle 0|v_2^\dagger v_2|0\rangle = 45696$.

References

- [1] W. Taylor, *M(atrix) Theory: Matrix Quantum Mechanics as a Fundamental Theory*, Rev. Mod. Phys. 73 (2001) 419-462, [arXiv:hep-th/0101126](#),
J. Hoppe, *Membranes and Matrix Models*, [arXiv:hep-th/0206192](#).
- [2] J. Wosiek, *On the SO(9) structure of supersymmetric Yang-Mills quantum mechanics* Phys. Lett. B619 (2005) 171-176, [arXiv:hep-th/0503236](#).
- [3] L. Brink, P. Ramond, X. Xiong, *Supersymmetry and Euler Multiplets*, JHEP10 (2002) 058, [arXiv:hep-th/0207253](#).
- [4] E. Cremmer, B. Julia and J. Scherk, *Supergravity theory in eleven dimensions*, Phys. Lett. B76, 409-412 (1978).

- [5] J. C. Plefka, A. Waldron, *On the quantum mechanics of M (atrix) theory*, Nucl. Phys. B512, 460-484 (1998), [arXiv:hep-th/9710104](#).
- [6] K. Millar, W. Taylor, M. Van Raamsdonk, *D-particle polarizations with multipole moments of higher-dimensional branes*, [arXiv:hep-th/0007157](#).
- [7] M. Bordemann, J. Hoppe, R. Suter, *Zero Energy States for $SU(N)$: A Simple Exercise in Group theory?*, [arXiv:hep-th/9909191](#).
- [8] R. Slansky, *Group theory for unified model building*, Phys. Rep., Vol. 79, Issue 1, p. 1-128 (1981).
- [9] M. Trzetrzelewski, *The number of gauge singlets in supersymmetric Yang-Mills quantum mechanics*, Phys. Rev. D76 (2007) 085012 , [arXiv:0708.2946](#).
- [10] Jin-Quan Chen, Jialun Ping and Fan Wang, *Group Representation Theory for Physicists*, World Scientific Publishing Company, Singapore (2002), 2nd Edition.
- [11] F.E. Browder, *Strongly elliptic systems of differential equations*, Annals of Mathematics Studies 33 (1954), 15-51.
- [12] D. Hasler, J. Hoppe, *Zero Energy States of Reduced Super Yang-Mills Theories in $d + 1 = 4, 6$ and 10 dimensions are necessarily $Spin(d)$ invariant*, [arXiv:hep-th/0211226](#).
- [13] J. Hoppe, *Asymptotic Zero Energy States for $SU(N)$ greater or equal 3*, [arXiv:hep-th/9912163](#).